

Effective Field Theory for Neutron-Deuteron Scattering: Higher Partial Waves

F. Gabbiani^a

*Department of Physics, Duke University
Box 90305, Durham, NC 27708-0305, USA
Email: fg@phy.duke.edu*

We present several theoretical results¹ for the phase shifts of higher partial waves for neutron-deuteron elastic scatterings in the quartet ($J = 3/2$) and doublet ($J = 1/2$) channels. This is the first case in which EFT has been successfully applied to these channels in a systematic way. Only the pionless case is investigated here, reserving the inclusion of pion exchanges for a later investigation.

Results for the quartet case are shown in Fig. 1. The phase shifts in degrees are plotted vs. the CM energy of the system. The only experimental results available in the literature are phase-shift analyses for processes with protons². Yet, their agreement with our theoretical predictions is excellent, indicating that electro-magnetic effects are not significant at the energies we study in this work. Potential models like the ones based on the Bonn and the av_{14} potentials³ give also similar results, but it is important to notice that we are able in principle not only to reproduce their results, but our effective theory framework with its well-defined power counting will in addition allow us to include higher order corrections and external currents in a consistent and systematic way.

The doublet case involves the solution of two coupled integral equations (see for example⁴). The S-wave component relates to the triton bound state, and will be studied in a future work. We show our results in Fig. 2.

1. F. Gabbiani, P. F. Bedaque and H. W. Griesshammer, Nucl. Phys. **A 675** (2000) 601.
2. E. Huttel *et al.*, Nucl. Phys. **A 406** (1983) 443; P. A. Schmelzbach *et al.*, Nucl. Phys. **A 197** (1972) 273.
3. A. Kievsky *et al.*, Nucl. Phys. **A 607** (1996) 402; A. Kievsky, private communication.
4. G. V. Skorniakov and K. A. Ter-Martirosian, Sov. Phys. **JETP 4** (1957) 648.

^aNew permanent address: Department of Physics and Astronomy, Iowa State University, Ames, IA 50011-3160




Figure 1: Real parts of the first four partial waves in the quartet channel of nd scattering versus the center-of-mass energy.




Figure 2: Real parts of the first four higher partial waves in the doublet channel of nd scattering versus the center-of-mass energy.